ELSEVIER

International Journal of Pharmaceutics 117 (1995) 257–259

international journal of pharmaceutics

Author Index

Akimoto, M., Furuya, A., Nakamura, M., Maki, T., Yamada, K., Suwa, T. and Ogata, H.
Release and absorption characteristics of chlorphenesin carbamate sustained-release formulations: in vitro-in vivo and in vivo dog-human correlations (117) 31
Alderborn, G., see Johansson, B. (117) 57
Alhaique, F., see Memoli, A. (117) 159
Arellano, A., see Santoyo, S. (117) 219
Arnardóttir, H.B., Sveinsson, S.J. and Kristmundsdóttir, T.
The use of a high intensity ultrasonic processor equipped with a flow cell in the production of reverse-phase liposomes (117) 237
Aungst, B.J. and Phang, S.
Metabolism of a neurotensin (8–13) analog by intestinal and nasal enzymes, and approaches to stabilize this peptide at these absorption sites (117) 95

Barry, B.W., see Cornwell, P.A. (117) 101
Bertoni, M., see Bonferoni, M.C. (117) 41
Bertoni, M., see Ferrari, F. (117) 49
Bonferoni, M.C., Rossi, S., Ferrari, F., Bertoni, M. and Caramella, C.
Influence of medium on dissolution-erosion behaviour of Na carboxymethylcellulose and on viscoelastic properties of gels (117) 41
Bonferoni, M.C., see Ferrari, F. (117) 49
Brandl, M., see Schneider, T. (117) 1
Brochart, H., see Goldbach, P. (117) 225
Buckton, G., Darcy, P. and Mackellar, A.J.
The use of isothermal microcalorimetry in the study of small degrees of amorphous content of powders (117) 253
Buckton, G., see Rillosi, M. (117) 75

Caramella, C., see Bonferoni, M.C. (117) 41
Caramella, C., see Ferrari, F. (117) 49
Cavalli, R., Morel, S., Gasco, M.R., Chetoni, P. and Saettone, M.F.
Preparation and evaluation in vitro of colloidal lipospheres containing pilocarpine as ion pair (117) 243
Chandler, S.G., Thomas, N.W. and Illum, L.
Nasal absorption in the rat: IV. Membrane activity of absorption enhancers. (117) 139
Chetoni, P., see Cavalli, R. (117) 243
Chiang, C.-C., Longer, M., Tyle, P., Fessler, D. and Shetty, B.
Formulation development of an oral dosage form for an HIV protease inhibitor, AG1284 (117) 197
Cornwell, P.A. and Barry, B.W.
Effects of penetration enhancer treatment on the statistical distribution of human skin permeabilities (117) 101
Cortesi, R., see Esposito, E. (117) 151

Darcy, P., see Buckton, G. (117) 253
Delgado-Charro, M.B. and Guy, R.H.
Iontophoretic delivery of nafarelin across the skin (117) 165

Ek, R., see Johansson, B. (117) 57
El-Gorashi, A.S., see Hosny, E.A. (117) 147
Esposito, E., Pastesini, C., Cortesi, R., Gambari, R., Menegatti, E. and Nastruzzi, C.
Controlled release of 1-β-D-arabinofuranosylcytosine (ara-C) from hydrophilic gelatin microspheres: in vitro studies (117) 151

Ferrari, F., Bertoni, M., Bonferoni, M.C., Rossi, S., Caramella, C. and Waring, M.J.
Comparative evaluation of hydrocolloid dressings by means of water uptake and swelling force measurements: II (117) 49
Ferrari, F., see Bonferoni, M.C. (117) 41
Fessler, D., see Chiang, C.-C. (117) 197
Furuya, A., see Akimoto, M. (117) 31

Gambari, R., see Esposito, E. (117) 151
Gasco, M.R., see Cavalli, R. (117) 243
Goldbach, P., Brochart, H., Wehrlé, P. and Stamm, A.
Sterile filtration of liposomes: retention of encapsulated carboxyfluorescein (117) 225
Gregoriadis, G., see Loukas, Y.L. (117) 85
Guy, R.H., see Delgado-Charro, M.B. (117) 165

Hanawa, T., see Watanabe, A. (117) 23
Hashida, M., see Yamashita, F. (117) 173
Hileman, G.A., see Katikaneni, P.R. (117) 13
Hino, T., see Schade, A. (117) 209

Hoffman, M., see Kedzierewicz, F. (117) 247
Hosny, E.A., Niazy, E.M. and El-Gorashi, A.S.
Effect of polycarbophil concentration on in vitro release and in vivo availability in beagle dogs of dihydroergotamine mesylate suppositories (117) 147
Hussain, M.A., Mersinger, L., Maurin, M.B. and Kettner, C.
In situ characterization of nasal leucine enkephalin degrading aminopeptidase. Susceptibility of the nasal enzyme to boronic acids and phosphorus-containing peptide and amino acid isosteres (117) 181

Illum, L., see Chandler, S.G. (117) 139

Jayasekera, P., see Loukas, Y.L. (117) 85
Johansson, B., Wikberg, M., Ek, R. and Alderborn, G.
Compression behaviour and compactability of microcrystalline cellulose pellets in relationship to their pore structure and mechanical properties (117) 57
Jones, D.S.
Preliminary investigations concerning the anti-adherence properties of polyhexamethylenebiguanide (Vantocil IB™) (117) 123

Kähkönen, K., see Mälkki-Laine, L. (117) 189
Katikaneni, P.R., Upadrashta, S.M., Rowlings, C.E., Neau, S.H. and Hileman, G.A.
Consolidation of ethylcellulose: effect of particle size, press speed, and lubricants (117) 13
Kawashima, Y., see Schade, A. (117) 209
Kedzierewicz, F., Villieras, F., Zinutti, C., Hoffman, M. and Maincent, P.
A 3 year stability study of tolbutamide solid dispersions and β-cyclodextrin complex (117) 247
Kettner, C., see Hussain, M.A. (117) 181
Kitano, M., see Yamashita, F. (117) 173
Koyama, Y., see Yamashita, F. (117) 173
Kristmundsdóttir, T., see Arnardóttir, H.B. (117) 237

Lanao, J.M., Vicente, M.T. and Sayalero, M.L.
Calculation of partial components of bioavailability in slow release formulations using model-independent methods (117) 113
Longer, M., see Chiang, C.-C. (117) 197
Loukas, Y.L., Jayasekera, P. and Gregoriadis, G.
Novel liposome-based multicomponent systems for the protection of photolabile agents (117) 85

Mackellar, A.J., see Buckton, G. (117) 253
Maincent, P., see Kedzierewicz, F. (117) 247
Maitani, Y., Yamamoto, T., Takayama, K. and Nagai, T.
The effect of soybean-derived sterol and its glucoside as an enhancer of nasal absorption of insulin in rabbits in vitro and in vivo (117) 129
Maki, T., see Akimoto, M. (117) 31
Mälkki-Laine, L., Purra, K., Kähkönen, K. and Tammilehto, S.
Decomposition of salbutamol in aqueous solutions. II. The effect of buffer species, pH, buffer concentration and antioxidants (117) 189
Martín, C., see Santoyo, S. (117) 219
Maurin, M.B., see Hussain, M.A. (117) 181
Memoli, A., Palermiti, L.G., Travagli, V. and Alhaique, F.
Egg and soya phospholipids – sonication and dialysis: A study on liposome characterization (117) 159
Menegatti, E., see Esposito, E. (117) 151
Mersinger, L., see Hussain, M.A. (117) 181
Morel, S., see Cavalli, R. (117) 243

Nagai, T., see Maitani, Y. (117) 129
Nakamura, M., see Akimoto, M. (117) 31
Nastruzzi, C., see Esposito, E. (117) 151
Neau, S.H., see Katikaneni, P.R. (117) 13
Niazy, E.M., see Hosny, E.A. (117) 147
Niwa, T., see Schade, A. (117) 209

Ogata, H., see Akimoto, M. (117) 31

Palermiti, L.G., see Memoli, A. (117) 159
Pastesini, C., see Esposito, E. (117) 151
Phang, S., see Aungst, B.J. (117) 95
Purra, K., see Mälkki-Laine, L. (117) 189

Rillosi, M. and Buckton, G.
Modelling mucoadhesion by use of surface energy terms obtained by the Lewis acid-Lewis base approach (117) 75
Rocco, W.L. and Swanson, J.R.
WIN 63843 polymorphs: prediction of enantiotropy (117) 231
Rößling, G., see Schneider, T. (117) 1
Rossi, S., see Bonferoni, M.C. (117) 41
Rossi, S., see Ferrari, F. (117) 49
Rowlings, C.E., see Katikaneni, P.R. (117) 13

Sachse, A., see Schneider, T. (117) 1
Saettone, M.F., see Cavalli, R. (117) 243
Santoyo, S., Arellano, A., Ygartua, P. and Martín, C.
Penetration enhancer effects on the in vitro percutaneous absorption of piroxicam through rat skin (117) 219
Sayalero, M.L., see Lanao, J.M. (117) 113
Schade, A., Niwa, T., Takeuchi, H., Hino, T. and Kawashima, Y.
Aqueous colloidal polymer dispersions of biodegradable DL-lactide/glycolide copolymer as basis for latex films: a new approach for the development of biodegradable depot systems (117) 209
Schneider, T., Sachse, A., Rößling, G. and Brandl, M.
Generation of contrast-carrying liposomes of defined size with a new continuous high pressure extrusion method (117) 1
Shetty, B., see Chiang, C.-C. (117) 197
Shively, M.L. and Thompson, D.C.
Oral bioavailability of vancomycin solid-state emulsions (117) 119
Stamm, A., see Goldbach, P. (117) 225

Sugihara, M., see Watanabe, A. (117) 23
Suwa, T., see Akimoto, M. (117) 31
Sveinsson, S.J., see Arnardóttir, H.B. (117) 237
Swanson, J.R., see Rocco, W.L. (117) 231

Takakura, Y., see Yamashita, F. (117) 173
Takayama, K., see Maitani, Y. (117) 129
Takeuchi, H., see Schade, A. (117) 209
Tammilehto, S., see Mälkki-Laine, L. (117) 189
Thomas, N.W., see Chandler, S.G. (117) 139
Thompson, D.C., see Shively, M.L. (117) 119
Travagli, V., see Memoli, A. (117) 159
Tyle, P., see Chiang, C.-C. (117) 197

Upadrashta, S.M., see Katikaneni, P.R. (117) 13

Vicente, M.T., see Lanao, J.M. (117) 113
Villieras, F., see Kedzierewicz, F. (117) 247

Waring, M.J., see Ferrari, F. (117) 49
Watanabe, A., Hanawa, T., Sugihara, M. and Yamamoto, K.
Development and pharmaceutical properties of a new oral dosage form of theophylline using sodium caseinate for the possible use in elderly patients (117) 23
Wehrlé, P., see Goldbach, P. (117) 225
Wikberg, M., see Johansson, B. (117) 57

Yamada, K., see Akimoto, M. (117) 31
Yamamoto, K., see Watanabe, A. (117) 23
Yamamoto, T., see Maitani, Y. (117) 129
Yamashita, F., Koyama, Y., Kitano, M., Takakura, Y. and Hashida, M.
Analysis of in vivo skin penetration enhancement by oleic acid based on a two-layer diffusion model with polar and nonpolar routes in the stratum corneum (117) 173
Ygartua, P., see Santoyo, S. (117) 219

Zinutti, C., see Kedzierewicz, F. (117) 247